

Original Research Article

Quercetin Suppresses Apoptotic Genes Expression and DNA Fragmentation Induced by Nano-rich Diesel Exhaust Particles in the Fetal Brain

¹Mohammed Eleyan, ¹Mohamed A. El-Desouky, ^{*2}Khairy A. Ibrahim and ³Hamed A. Daboun

Abstract

¹Biochemistry Division, Faculty of Science, Cairo University, Egypt.

²Central Agricultural Pesticides Laboratory, Agricultural Research Center, Dokki, Giza, Egypt.

³Department of Chemistry, Faculty of Science, Cairo University, Egypt.

*Corresponding Author's E-mail: khairy_moneim@yahoo.com

Nano-rich diesel exhaust particles (NRDEPs) are an extensive range of toxic metals existing generally on environmental system and associated with nervous system disorders, including autism spectrum and induce apoptosis through oxidative stress. Our aim was to evaluate the potential effects of NRDEPs on the brain tissue as apoptosis inducer agent via oxidative stress in hand and the quercetin as powerful anti-oxidant, anti-apoptotic on other hand. Forty pregnant (from the 5th to the 20th day of gestation) albino rats were randomly divided into control group which received normal saline and quercetin group which received 100 mg/kg quercetin. The 3rd one received 20 mg/kg of NRDEPs and the 4th received 100 mg/kg quercetin two hours prior to NRDEPs exposure. The dams were anaesthetized by the end of gestation and fetal brain was removed for immunohistochemistry, biochemical parameters estimation, comet assay and real time polymerase chain reaction (RT-PCR). Our results showed that maternally co-treatment with quercetin diminished NRDEPs-induced oxidative stress (reduced MDA level, increased GSH level and enhanced both GST and PON activities). Moreover, quercetin also down regulated the expression of Bax and caspase-3 and reduced DNA fragmentation. These outcomes indicate that quercetin has a suppressive role against oxidative stress and apoptosis induced by nano-rich diesel exhaust particles in rat fetal brain.

Keywords: Diesel exhaust, Nanoparticles, Quercetin, Apoptosis, Fetus.

Abbreviations

Bax, Bcl-2-associated X protein; Bcl-2, B-cell lymphoma 2; Bcl-XL, B-cell lymphoma-extra large; CNTs, carbon nanotubes; DE, diesel exhaust; NP, nanoparticles; NRDEPs, nano-rich diesel exhaust particles; PAHs, Polycyclic Aromatic Hydrocarbons; PM, Particulate Matter; ROS, Reactive oxygen species.

INTRODUCTION

Diesel exhaust contains a wide range of organic and inorganic compounds that have been described as

particulate matter air pollutants (Kagawa, 2002). As diesel fuel undergoes combustion in engines, it produces

particles known as nano-rich diesel exhaust particles (NRDEPs) which composes of some different chemicals with different sizes and physical proprieties such as polycyclic aromatic hydrocarbons, toxic metals, diesel derived nanoparticles (NPs), carbon nanotubes (CNTs) and carbon nanofibers (Jung et al., 2013).

NRDEPs are very toxic because of their minute size, large reactive surface area and modified quantum effects (Buzea et al., 2007). Once inhaled, NRDEPs can cross the cellular barriers, transfer through the blood circulation and finally accumulate in secondary organs as brain, liver, heart and spleen (Semmler-Behnke et al., 2014).

Previous studies have shown that the trophoblastic cells and fetal capillaries relative volume decreased after maternal exposure to NRDEPs as they have ability to pass through the placenta as well as reach the blood stream of embryo (Valentino et al., 2016; Veras et al., 2008). NRDEPs can affect the fetal central nervous system development which leads to pathological abnormalities of fetal brain and various neurological disorders (Sugamata et al., 2006).

Recently, the potential relationship between oxidative stress and many diseases pathophysiology in human and animal has been getting an inordinate deal of attention. Oxidative stress induced by nanoparticles has been known as a source of developmental toxicity (Sun et al., 2013). NRDEPs induce oxidative stress by generation of hydroxyl and oxygen free radicals along with reduction in the cellular antioxidant defenses system (Han et al., 2001; Ross and Kasum, 2002). NRDEPs is also reported to generate reactive oxygen species and induce the programmed cell death (Hiura et al., 2000).

Neuroprotective approaches support the supplementation of a valuable, highly efficient and low-cost of natural antioxidants which maintains the balance between oxidant and cellular antioxidant in turn protect the central nerves system from oxidative stress-related diseases, neural degeneration and apoptosis. Quercetin is a bioactive antioxidant plant that has an extensive range of health benefits and distributes over a large number of fruits, vegetables and medical herbs (Slikkerm et al., 1999). Quercetin also has various protective and ameliorative functions including, metal chelator, free radical's scavenger so it may prevent oxidative stress, cell injury and apoptosis (Bu et al., 2012).

Although embryonic organs are more sensitive to toxic materials than adults and the data about the suppressive role of quercetin against apoptosis induced by NRDEPs in fetal brain is limited, so evaluating the safety of nanoparticles is very important at this critical period of pregnancy. Consequently, the present aim was to divulge the association between oxidative stress and NRDEPs-induced apoptosis on the brain of albino rat fetuses in hand and the suppressive role of quercetin on the other hand.

MATERIALS AND METHODS

Chemicals used

The NRDEPs were obtained from the Egyptian Transport Network. Quercetin (2,3',4',5,7-pentahydroxyflavone), other chemicals and substrates were obtained from Sigma Chemical Company (U.S.A.).

Experimental design

Forty pregnant albino rats (*Rattus norvegicus*) were randomly divided into four experimental groups as follows: control group (G1): dams received normal saline, quercetin group (G2): received 100 mg/kg quercetin, NRDEPs group (G3): received 20 mg/kg of NRDEPs with intra-tracheal instillation, quercetin + NRDEPs group (G4): received 100 mg/kg quercetin two hours prior to the intra-tracheal instillation of 20 mg/kg NRDEPs. The experimental study was done from the 5th to the 20th day of gestation, and this protocol was approved by The Institutional Animal Care and Use Committee (IACUC) at Cairo University (CU.1.S.74.17).

Specimen

By the end of gestation, the dams were anaesthetized and brain specimens were removed immediately from the fetus, washed, dried and fixed by formalin for immunohistochemistry. Other fatal brain tissues were kept frozen at -80°C for biochemical analysis, comet assay and real-time polymerase chain reaction (Real-Time PCR).

Biochemical analysis

Preparation of the tissue fraction

Brain tissues were homogenized in pre cold saline, then the homogenate was centrifuged and the resultant supernatant was used for MDA and GSH levels measurement as well as PON and GST activities assessment. The protein concentration was measured calorimetrically by the method adopted by (Bradford 1976) with bovine serum albumin as standard using Spectronic Helios Alpha UV-Vis (England).

Determination of paraoxonase (PON) activity

The activity of paraoxonase was determined spectrophotometrically and p-nitrophenyl acetate was employed as a substrate, as previously described (Gan et al., 1991). The enzyme activity was measured by the

Table 1. Oligonucleotide primers used in SYBR Green RT-PCR

Gene	Primer sequence (5'-3')	Accession Number	Reference
β-actin	F-TCCTCCTGAGCGCAAGTACTCT	V01217	(Banni et al., 2010)
	R-GCTCAGTAACAGTCCGCCTAGAA		
BAX	F-CACCAGCTCTGAACAGATCATGA	RRU49729	(Kinouchi 2003)
	R-TCAGCCCATCTTCTCCAGATGGT		
Caspase-3	F-AGTTGGACCCACCTTGTGAG	NM_012922.2.298	(Shi et al., 2009)
	R-AGTCTGCAGCTCCTCCACAT		

increase in the developed yellow colour through p-nitrophenol released.

Determination of glutathione-S-transferase (GST) activity

The activity of GST was assayed spectrophotometrically following a previously published method (Habig et al., 1974). This method based on measure the yellow colour developed as a result of the conjugation of 1-chloro-2, 4-dinitrobenzene (CDNB) with reduced glutathione (Habdous et al., 2002).

Determination of reduced glutathione (GSH) content

Reduced glutathione content was determined as previously described (Ellman 1959). The method is based on the determination of a yellow color that develops when [5, 5 dithiol- bis (2-dinitrobenzoic acid)] (DTNB) is added to sulphhydryl compounds, the developed color was measured spectrophotometrically at 412 nm.

Determination of lipid peroxidation level

The method is based on the determination of MDA as an end product of lipid peroxidation, which can react with thiobarbituric acid in acidic medium to yield a pink colored of trimethine complex exhibiting an absorption maximum at 532 nm (Yoshioka et al., 1979).

Immunohistochemistry

Paraffin embedded brain tissues were cut into five μm sections to detect caspase-3 and Bax proteins. Rabbit caspase-3 and Bax antibodies were used as a primary antibody. Formerly biotinylated secondary antibody was added, and then horse-radish peroxidase conjugated with streptavidin. As streptavidin has a high affinity toward biotin, it binds to the site where primary antibody coated the background, after adding a chromagen (DAB) a reddish color appears (Sati et al., 2010), a light microscope with a digital camera were used for images capture.

Comet assay


The comet assay was applied under alkaline conditions using ordinary microscope slides as previously described by (Collins 2004; Singh et al., 1988). The brain tissues of fetuses were minced in Hank's balanced salt solution, (HBSS) pH 6.5 (Ca⁺⁺ and Mg⁺⁺ free) with 20 mM EDTA and 10% DMSO). Minced brain tissues were collected and centrifuged at 2,000 rpm at 4 C for 10 min. cell pellet was suspended in PBS-CMF. The slides (were prepared one day before) were pre-coated with 200 μL of 1.5% normal melting point agarose and dried up at room temperature. Then a mixture of 10 μL of cell suspension and 75 μL of 0.5% low melting point agarose was added. The covers were gently slid off and third layer (80 μL of 1.0% low melting point agarose) was added to the slides. The covers were replaced and the slides were allowed to harden for 3–5 minutes and the slides were slowly immersed in lysing solution (2.5 M NaCl, 100 mM EDTA, 10 mM Trizma base with pH 10. Freshly 1% Triton X-100 and 10% DMSO) and refrigerated for one hour. The slides were gently removed from lysing solution and rinsed carefully to remove detergents and salts. The slides were placed close together on horizontal gel electrophoresis (25 V and 300 mA for 40 min; JSB-30, USA). After the electrophoresis, slides were gently washed three times using neutralization buffer. Then 70 % ethanol was used for fixing the slides for 10 min and dried up at room temperature. Then the slides were stained with aqueous ethidium bromide solution. Fluorescence microscope (Lx 400, Labomed, USA) was used for DNA fragmentation detection. The degree of DNA damage was determined by v1.5 of Comet Score.

Expression of apoptosis genes determination

RNA was extracted from fetal brain according to RNeasy Mini Kit instructions (Catalogue no.74104). RNA was then reverse transcribed using (Revert Aid Reverse Transcriptase Thermo Fisher, Catalog number: K1622) according to manufacturer's instructions. Primers were obtained from (Metabion, Germany) and the sequences of these primers obtainable in Table 1. Real-time quantification was performed in the Stratagene MX3005P instrument using the Quantitect SYBR green PCR kit (Cat. No. 204141). The reactions were performed in a

Table 2. Effects of NRDEPs and co-treatment with quercetin on MDA and GSH levels, GST and PON activities in fetal brain tissues

	MDA (nmol/g tissue)	GSH (μ M/g tissue)	GST (μ M/min/mg protein)	PON (μ M/min/mg protein)
G1	10.02 \pm 0.6	20.16 \pm 0.67	524.55 \pm 20.49	123.9 \pm 3.93
G2	9.688 \pm 0.44 ^{c,d}	21.87 \pm 0.84 ^{c,d}	517.49 \pm 10.08 ^c	125.02 \pm 2.73 ^{c,d}
G3	21.68 \pm 0.82 ^{a,b,d}	15.15 \pm 0.72 ^{a,b,d}	372.02 \pm 8.59 ^{a,b,d}	89.56 \pm 3.32 ^{a,b,d}
G4	15.7 \pm 0.55 ^{a,b,c}	18.61 \pm 0.57 ^{b,c}	476.19 \pm 16.07 ^c	109.01 \pm 5.02 ^{a,b,c}

**Figure 1.** A photomicrograph of caspase-3 (top) and Bax (bottom) immunohistochemical staining of fetal brain sections on the 20th day of gestation; (A,E) control group; (B,F) quercetin group; (C,G) NRDEPs group; (D,H) NRDEPs + quercetin group.(Magnification \times 40)

25 μ l volume mix containing; 12.5 μ l 2x SYBR Green PCR Master Mix, 1 μ l primers, 2 μ l cDNA, and 8.5 μ l of RNase Free Water. Dissociation stage was performed after the amplification to verify the specificity of the PCR products.

Statistical analysis

Data are presented as mean \pm standard error of mean (SEM). Differences between groups were achieved by the analysis of variance (ANOVA) followed by Tukey's multiple comparison post hoc analysis. Significance was set at probability $P \leq 0.05$ using IBM SPSS version 22 software package (SPSS, IBM, Chicago, IL, USA).

RESULTS

Malondialdehyde (MDA) level

The biochemical results presented in this study revealed that fetal brain MDA level was significantly increased ($P < 0.05$) in both NRDEPs and NRDEPs+ quercetin groups as compared with control one on the other hand the co-treatment with quercetin significantly ameliorated the fetal brain MDA level (Table 2).

Antioxidant system of fetal brain

Maternally intoxication with 20 mg/kg NRDEPs pointedly disrupted the antioxidant system of fetal brain as GSH level as well as GST and PON activities were significantly decreased ($P < 0.05$) as compared with control. Inside out, the co-treatment with quercetin exhibited a significant increase in GSH level ($P < 0.05$) and enhanced the GST and PON activities as compared with the NRDEPs group (Table 2).

Each value represents the mean and SME. One-way analysis of variance (ANOVA) followed by Turkey's honestly significant difference (HSD) test, the $p \leq 0.05$ level was set as statistically significant different; (a) significant compared to control, (b) significant compared to quercetin group, (c) significant compared to NRDEPs group, (d) significant compared to NRDEPs + quercetin group.

Results of Caspase-3 and Bax Transcripts Using Real-Time RT-PCR

The Bax and caspase-3 genes expression exhibited significantly up-regulated ($P < 0.05$) by 22.4 and 28.7 folds respectively, among NRDEPs group and by 2.7 and 3.3 folds respectively, among NRDEPs + quercetin group




Figure 2. Fold changes of caspase-3 and Bax genes expression in brain of albino rat fetuses maternally exposed to the NRDEPs and co-treatment with quercetin; (A) Bax gene and (B) caspase-3 gene.




Figure 3. A photomicrograph of comet assay in fetal brain (A) control group; (B) quercetin group; (C) NRDEPs group; (D) NRDEPs + quercetin group.

Table 2. Effects of NRDEPs and co-treatment with quercetin on comet parameters in fetal brain tissues

	Head DNA (%)	Tail DNA (%)	Tail Length (μm)	Tail Moment (AU)
G1	91.65 ± 1.78	8.35 ± 1.67	3.13 ± 0.025	0.535 ± 0.036
G2	90.45 ± 1.94 ^{c,d}	9.55 ± 1.92 ^{c,d}	3.46 ± 0.22 ^{c,d}	0.578 ± 0.015 ^{c,d}
G3	77.68 ± 1.76 ^{a,b,d}	22.32 ± 1.75 ^{a,b,d}	5.23 ± 0.25 ^{a,b,d}	1.35 ± 0.177 ^{a,b,d}
G4	82.7 ± 1.19 ^{a,b,c}	17.3 ± 1.19 ^{a,b,c}	3.58 ± 0.06 ^{a,b,c}	0.7 ± 0.072 ^{a,b,c}

as both compared with control. However, non-significant down regulation ($P < 0.05$) was observed between quercetin and control groups. Inside out, the co-treatment with quercetin revealed an efficient down-regulation of both Bax and caspase-3 gene expression in fetal brain tissues, shown in (Fig. 2A and 2B).

Immunohistochemistry

The brain tissues of the NRDEPs group revealed a positive reaction for active caspase-3 (Fig.1 C) and Bax (Fig.1 G) in the neurons as compared with control (Fig.1 A and E) and quercetin (Fig.1 B and F) and groups that showed negative reaction for caspase-3 and Bax. Quercetin co-treatment attenuated the up-regulation of caspase-3 (Fig.1 D) and Bax (Fig.1 H).

Fetal brain DNA damage

In this study, comet assay demonstrated the genotoxicity of NRDEPs as evidenced by elevation of all estimated parameters of DNA damage (% DNA in tail, % DNA in head, Tail length and Tail moment) in fetal brain tissues compared to control values. On the other hand the co-treatment with quercetin significantly minimized the DNA damage induced by NRDEPs as shown in (tables 3 and Fig. 3).

Each value represents the mean and SME. One-way analysis of variance (ANOVA) followed by Turkey's honestly significant difference (HSD) test, the $p \leq 0.05$ level was set as statistically significant different. (a) significant compared to control, (b) significant compared to quercetin group, (c) significant compared to NRDEPs group, (d) significant compared to

NRDEPs + quercetin group.

DISCUSSION

Nano-rich diesel exhaust particles are worldwide relevant air pollutants that exert a fierce human and animal health effect (Dockery et al., 1993), so our study was conducted to evaluate the suppressive role of quercetin against apoptosis induced by nano-rich diesel exhaust particles in albino rat fetal brain.

NRDEPs induce oxidative stress by (I) transition element-based nanoparticles such as CNT (II) free radicals present on surfaces of particles such as quartz and carbonaceous particles, (III) redox-active sites resulting from functionalization of nanoparticles as quinones capable of generating ROS (Knaapen et al., 2004; Li et al., 2010) (IV) NRDEPs chemical composition such as transition metals (Al, Fe, Mn, Cu, Pb, Zn, Cr, V, Ba and Ni) as well as PAH and their derivatives (Andersson et al., 1998). The distinctive physicochemical structures of nanoparticles, as their nanosize with large reactive surface area are responsible for generating ROS and oxidative injury (Auffan et al., 2009). Beside their direct or indirect damaging effects on cellular lipids, proteins, and DNA, an increased production of ROS triggers programmed cell death.

Surely, NRDEPs pass through the placenta, then taken by the fetuses and entered fetal circulation and reach the brain after passing through the not fully developed fetal blood brain barrier which lead to induce oxidative stress and apoptosis (Hougaard et al., 2008; Sugamata et al., 2006).

Elevated levels of MDA in the current study, revealed a damaging effects on cell membrane structure and functions by the theory of lipid peroxidation. Oxidative stress provoked by free radicals such as H_2O_2 is stated to be a vital mechanism of many infections and is currently evaluated to be one of the main reasons leading to the adverse health effects induced by NRDEPs. ROS production and changes in cell cycle metabolism contributing to oxidative stress in NRDEPs exposed cells (Cachon et al., 2014).

Quercetin showed an effective antioxidant property that is most often attributed to its hydroxyl groups, which relates with its electron donating capability (Tu et al., 2015). It is supposed that quercetin prevent the free radical chain reaction progression through the mechanism of free radicals trapping at the interface of the membranes (Ross and Kasum 2002).

According to our results, co-treatment with quercetin after 2 hours of exposure to NRDEPs attenuated lipid peroxidation due to its ability to inhibit lipid peroxidation (Bakheet 2011), quercetin is an effective ions chelator and can chelate transition metal ions present in NRDEPs that may responsible for increasing oxidative stress (Rice-Evans et al., 1996).

The drop in GSH level may due to scavenging of NRDEPs – induced free radicals production, suppression GSH synthesis and/or restriction of intracellular reduction of oxidized GSH (Mohan et al., 2013).

The decrease in GST activity might be owing to GST participated in detoxifying NRDEPs to non-toxic metabolites or by rapidly binding and very slowly turning over these xenobiotics (Jebur et al., 2014; Lonare et al., 2014). Our results suggest that NRDEPs or its metabolites, other than free radicals, were responsible for the reduction in PON activity which may act as scavenger for these radicals. NRDEPs contain a variety of reactive components that are highly reactive towards thiols forming sulfhydryl-ether linkage (Nishio and Watanabe 1997), and/or able to connect with Zn^{2+} ion in the active center of PON enzyme and it can diminish the PON activity (Demir 2011). Otherwise NRDEPs might have negative effects on PON activities or on their 3-dimensional structures that result in some changes in the metabolisms (CEBECI et al., 2014).

The improved GSH, GST and PON activities propose that quercetin mediated the modulation of cellular antioxidant levels. In agree with several previous studies which reported that quercetin can elevate GSH, GST and PON as well as reduce lipid peroxidation levels (Attia 2010; Kapiszewska et al., 2007; Papiez et al., 2008).

Our immunohistochemistry and RT-PCR results showed that the expression level of Bax and caspase-3 in fetal brain were significantly up-regulated after maternally intoxicated with NRDEPs.

We suggested that the NRDEPs may produce free radical and induce oxidative stress as result from inhibition of the enzymatic and nonenzymatic defenses systems. If this suggestion is valid, NRDEPs may induce apoptosis via mitochondrial intrinsic pathway through activation of caspase-9 which turn on activation of caspase-3 that triggers the rapid DNA fragmentation.

In fact, there is a very strong relation between high oxidative stress and apoptosis. Apoptosis has two pathways: the intrinsic and the extrinsic pathways. The intrinsic pathway implicates mitochondrial perturbation causing cytotoxicity (Danial and Korsmeyer 2004; Marnett 1999; Sun et al., 1999).

The Bcl-2 family consists of proteins that have anti-apoptotic (e.g., Bcl-2 and Bcl-XL) and a pro-apoptotic (e.g., BH3 and Bax) members, which exerts an important role in the controlling of intrinsic mitochondrial pathway of programmed cell death (Desagher et al., 1999). High Bax mRNA expression levels lead to formation of homo- or heterodimers by its ability to interacting with other Bcl-2 members resulted in permeabilization of the outer membrane of mitochondrial (Szabo and Zoratti 2014) which causes disturbance in its transmembrane potential (Kroemer et al., 2007) This lead to release of cytochrome c from mitochondria into the cytoplasm and formation of a complex known as apoptosome (Lü et al., 2003).

Apoptosome can activate caspase-9 that sequentially activates the effector caspase-3 (Johnson and Jarvis 2004). Caspase-3 is responsible for destruction of the specific proteins that lead to initiates apoptosis by releasing caspase-activated deoxyribonuclease which triggers rapid DNA fragmentation (Han et al., 2011) as described in our study by comet assay.

It was also reported that NRDEP induced oxidative stress via production of reactive oxygen species which leads to mitochondrial toxicity or reduce mitochondrial activity, induce mitochondrial O₂ – generation (Hiura et al., 2000; Tseng et al., 2015), which are highly reactive with DNA and induce both apoptosis and DNA fragmentation (Gustafsson and Gottlieb, 2007; Shen et al., 2013).

In the present study, quercetin co-treatment effectively down-regulated the high expression of caspase-3 and Bax, which they are often revealing the degree of apoptosis in fetal brain after maternally exposure to NRDEPs.

Quercetin has antiapoptotic function, which has effectively suppressed of Bax-dependent apoptosis and inhibited of several proapoptotic transcription factors along with up-regulation of anti-apoptotic member Bcl-XL (Attia 2010; Bu et al., 2012).

CONCLUSION

On conclusion, nano-rich diesel exhaust particles induce up-regulation of Bax and caspase-3 level expression leading to trigger the intrinsic mitochondrial apoptosis pathway through oxidative stress in addition to DNA fragmentation. Nevertheless the potent antioxidant quercetin has ameliorative roles against NRDEPs induced oxidative stress along with enhancing cellular antioxidant defense mechanism and inhibiting Bax and caspase-3 level expression so, it can alleviate apoptosis and DNA fragmentation.

Conflict of interest

The authors declare that they have no competing of interest.

Ethics approval and consent to participate

See MATERIALS AND METHODS.

REFERENCES

Andersson H, Lindqvist E, Westerholm R, Grägg K, Almén J, Olson L (1998). Neurotoxic effects of fractionated diesel exhausts following microinjections in rat hippocampus and striatum. *Environmental research* 76(1):41-51.

Attia SM (2010). The impact of quercetin on cisplatin-induced clastogenesis and apoptosis in murine marrow cells. *Mutagenesis* 25(3):281-288.

Auffan M, Rose J, Bottero J-Y, Lowry GV, Jolivet J-P, Wiesner MR (2009). Towards a definition of inorganic nanoparticles from an environmental, health and safety perspective. *Nature nanotechnology* 4(10):634.

Bakheet SA (2011). Assessment of anti-cytogenotoxic effects of quercetin in animals treated with topotecan. *Oxidative medicine and cellular longevity* 2011; 2011: 824597.

Banni M, Messaoudi I, Said L, El Heni J, Kerkeni A, Said K (2010). Metallothionein gene expression in liver of rats exposed to cadmium and supplemented with zinc and selenium. *Archives of environmental contamination and toxicology* 59(3):513-519.

Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry* 72(1-2):248-254.

Bu T-l, Jia Y-d, Lin J-x, Mi Y-l, Zhang C-q (2012). Alleviative effect of quercetin on germ cells intoxicated by 3-methyl-4-nitrophenol from diesel exhaust particles. *J. Zhejiang Uni. Sci. B* 13(4):318-326.

Buzaa C, Pacheco II, Robbie K (2007). Nanomaterials and nanoparticles: Sources and toxicity. *Biointerphases* 2(4):MR17-MR71.

Cachon BF, Firmin S, Verdin A, Ayi-Fanou, L, Billet S, Cazier F, Martin PJ, Aissi F, Courcot D, Sanni A (2014). Proinflammatory effects and oxidative stress within human bronchial epithelial cells exposed to atmospheric particulate matter (PM_{2.5} and PM_{>2.5}) collected from Cotonou, Benin. *Environmental pollution* 185:340-351

Cebeci B, Alim Z, Beydemir Ş (2014). In vitro effects of pesticide exposure on the activity of the paraoxonase-1 enzyme from sheep liver microsomes. *Turkish J. Chem.* 38(3):512-520

Collins AR (2004). The comet assay for DNA damage and repair. *Molecular biotechnology* 26(3):249

Danial NN, Korsmeyer SJ (2004). Cell Death: Critical Control Points. *Cell* 116(2):205-219 doi:https://doi.org/10.1016/S0092-8674(04)00046-7

Demir N (2011). An, In Vitro, Study of some Pesticides on the Activity of Human Serum Paraoxonase (PON1).

Desagher S, Osen-Sand A, Nichols A, Eskes R, Montessuit S, Lauper S, Maundrell K, Antonsson B, Martinou JC (1999) Bid-induced Conformational Change of Bax Is Responsible for Mitochondrial Cytochrome c Release during Apoptosis. *The Jo. Cell Biol.* 144(5):891-901.

Dockery DW, Pope CA, Xu X, Spengler JD, Ware JH, Fay ME, Ferris JBG, Speizer FE (1993) An association between air pollution and mortality in six US cities. *New Eng. J. Med.* 329(24):1753-1759.

Ellman GL (1959). Tissue sulfhydryl groups. *Archives of biochemistry and biophysics* 82(1):70-77

Gan KN, Smolen A, Eckerson HW, La Du BN (1991). Purification of human serum paraoxonase/arylesterase. Evidence for one esterase catalyzing both activities. *Drug Metabolism and Disposition* 19(1):100-106.

Gustafsson ÅB, Gottlieb RA (2007). Bcl-2 family members and apoptosis, taken to heart. *American Journal of Physiology-Cell Physiology* 292(1):C45-C51.

Habdous M, Vincent-Viry M, Visvikis S, Siest G (2002). Rapid spectrophotometric method for serum glutathione S-transferases activity. *Clinica Chimica Acta* 326(1-2):131-142.

Habig WH, Pabst MJ, Jakoby WB (1974). Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *Journal of biological Chemistry* 249(22):7130-7139 .

Han B, Wang Q, Cui G, Shen X, Zhu Z (2011). Post-treatment of Bax-inhibiting peptide reduces neuronal death and behavioral deficits following global cerebral ischemia. *Neurochemistry International* 58(2):224-233.

Han J-Y, Takeshita K, Utsumi H (2001). Noninvasive detection of hydroxyl radical generation in lung by diesel exhaust particles. *Free Radical Biology and Medicine* 30(5):516-525.

Hiura TS, Li N, Kaplan R, Horwitz M, Seagrave J-C, Nel AE (2000). The role of a mitochondrial pathway in the induction of apoptosis by chemicals extracted from diesel exhaust particles. *The J. Immunol.* 165(5):2703-2711.

- Hougaard KS, Jensen KA, Nordly P, Taxvig C, Vogel U, Saber AT, Wallin H (2008) Effects of prenatal exposure to diesel exhaust particles on postnatal development, behavior, genotoxicity and inflammation in mice. *Particle and fibre toxicology* 5(1):3.
- Jebur AB, Nasr HM, El-Demerdash FM (2014). Selenium modulates β -cyfluthrin-induced liver oxidative toxicity in rats. *Environmental toxicology* 29(11):1323-1329.
- Johnson CR, Jarvis WD (2004). Caspase-9 regulation: An update. *Apoptosis* 9(4):423-427.
- Jung HS, Miller A, Park K, Kittelson DB (2013). Carbon nanotubes among diesel exhaust particles: real samples or contaminants? *Journal of the Air & Waste Management Association* 63(10):1199-1204.
- Kagawa J (2002). Health effects of diesel exhaust emissions—a mixture of air pollutants of worldwide concern. *Toxicology* 181-182:349-353.
- Kapiszewska M, Cierniak A, Papiez MA, Pietrzycka A, Stepniewski M, Lomnicki A (2007). Prolonged Quercetin Administration Diminishes the Etoposide-Induced DNA Damage in Bone Marrow Cells of Rats. *Drug and Chemical Toxicology* 30(1):67-81.
- Kinouchi S (2003). Changes in apoptosis-related genes (Bcl-2, Bax) in the urethras of old female rats following estrogen replacement. *Yonago Acta Medica* 46:109-115.
- Knaapen AM, Borm PJ, Albrecht C, Schins RP (2004). Inhaled particles and lung cancer. Part A: Mechanisms. *International Journal of Cancer* 109(6):799-809.
- Kroemer G, Galluzzi L, Brenner C (2007). Mitochondrial Membrane Permeabilization in Cell Death. *Physiological Reviews* 87(1):99-163 doi:10.1152/physrev.00013.2006.
- Li J-Je, Muralikrishnan S, Ng C-T, Yung L-YL, Bay B-H (2010). Nanoparticle-induced pulmonary toxicity. *Experimental biology and medicine* 235(9):1025-1033.
- Lonare M, Kumar M, Raut S, Badgujar P, Doltade S, Telang A (2014). Evaluation of imidacloprid-induced neurotoxicity in male rats: A protective effect of curcumin. *Neurochemistry international* 78:122-129.
- Lü C-X, Fan T-J, Hu G-B, Cong R-S (2003). Apoptosis-inducing factor and apoptosis. *Sheng Wu Hua Xue Yu Sheng Wu Wu Li Xue Bao (Shanghai)* 35(10):881-885.
- Marnett LJ (1999). Lipid peroxidation—DNA damage by malondialdehyde. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 424(1):83-95.
- Mohan D, Thiyagarajan D, Murthy PB (2013). Toxicity of exhaust nanoparticles. *African Journal of Pharmacy and Pharmacology* 7(7):318-331.
- Nishio E, Watanabe Y (1997). Cigarette smoke extract inhibits plasma paraoxonase activity by modification of the enzyme's free thiols. *Biochemical and biophysical research communications* 236(2):289-293.
- Papiez MA, Cierniak A, Krzysciak W, Bzowska M, Taha HM, Jozkowicz A, Piskula M (2008). The changes of antioxidant defense system caused by quercetin administration do not lead to DNA damage and apoptosis in the spleen and bone marrow cells of rats. *Food and Chemical Toxicology* 46(9):3053-3058.
- Rice-Evans CA, Miller NJ, Paganga G (1996). Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free radical biology and medicine* 20(7):933-956.
- Ross JA, Kasum CM (2002). Dietary flavonoids: bioavailability, metabolic effects, and safety. *Annual review of nutrition* 22(1):19-34.
- Sati L, Seval-Celik Y, Demir R (2010). Lung surfactant proteins in the early human placenta. *Histochemistry and cell biology* 133(1):85.
- Semmler-Behnke M, Lipka J, Wenk A, Hirn S, Schäffler M, Tian F, Schmid G, Oberdörster G, Kreyling WG (2014). Size dependent translocation and fetal accumulation of gold nanoparticles from maternal blood in the rat. *Particle and Fibre Toxicology* 11:33-33.
- Shen B, He P-J, Shao C-L (2013). Norcantharidin induced DU145 cell apoptosis through ROS-mediated mitochondrial dysfunction and energy depletion. *PLoS One* 8(12):e84610.
- Shi Y, Song Y, Wang Y, Liang X, Hu Y, Guan X, Cheng J, Yang K (2009). p, p'-DDE induces apoptosis of rat Sertoli cells via a FasL-dependent pathway. *J Biomed Biotechnol*, 2009; 181282.
- Singh NP, McCoy MT, Tice RR, Schneider EL (1988). A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental cell research* 175(1):184-191.
- Slikkerm W, Youdim C, Hallc P, Tremblay W (1999) The Future of Neuroprotection. *Annals of the New York Academy of Sciences* 890(1):529-533.
- Sugamata M, Ihara T, Takano H, Oshio S, Takeda K (2006). Maternal diesel exhaust exposure damages newborn murine brains. *Journal of health science* 52(1):82-84.
- Sun J, Zhang Q, Wang Z, Yan B (2013) Effects of nanotoxicity on female reproductivity and fetal development in animal models. *International journal of molecular sciences* 14(5):9319-9337.
- Sun X-M, MacFarlane M, Zhuang J, Wolf BB, Green DR, Cohen GM (1999). Distinct Caspase Cascades Are Initiated in Receptor-mediated and Chemical-induced Apoptosis. *Journal of Biological Chemistry* 274(8):5053-5060.
- Szabo I, Zoratti M (2014) Mitochondrial Channels: Ion Fluxes and More. *Physiological Reviews* 94(2):519-608.
- Tseng C-Y, Wang J-S, Chang Y-J, Chang J-F, Chao M-W (2015). Exposure to High-Dose Diesel Exhaust Particles Induces Intracellular Oxidative Stress and Causes Endothelial Apoptosis in Cultured In Vitro Capillary Tube Cells. *Cardiovascular Toxicology* 15(4):345-354.
- Tu B, Liu Z-J, Chen Z-F, Ouyang Y, Hu Y-J (2015). Understanding the structure–activity relationship between quercetin and naringenin: in vitro. *RSC Advances* 5(128):106171-106181
- Valentino SA, Tarrade A, Aioun J, Mourier E, Richard C, Dahirel M, Rousseau-Ralliard D, Fournier N, Aubrière MC, Lallemand MS, Camous S, Guinot M, Charlier M, Aujean E, Al Adhami H, Fokkens PH, Agier L, Boere JA, Cassee FR, Slama R, Chavatte-Palmer P (2016). Maternal exposure to diluted diesel engine exhaust alters placental function and induces intergenerational effects in rabbits. *Particle and Fibre Toxicology* 13(1):39.
- Veras MM, Damaceno-Rodrigues NR, Caldini EG, Ribeiro A.ACM, Mayhew TM, Saldiva PHN, Dolnikoff M (2008). Particulate Urban Air Pollution Affects the Functional Morphology of Mouse Placenta1. *Biology of Reproduction* 79(3):578-584.
- Yoshioka T, Kawada K, Shimada T, Mori M (1979). Lipid peroxidation in maternal and cord blood and protective mechanism against activated-oxygen toxicity in the blood. *American journal of obstetrics and gynecology* 135(3):372-376.